

[2008]

Gretarsdottir S, Thorleifsson G, Manolescu A, Styrkarsdottir U, Helgadottir A, Gschwendtner A, Kostulas K, Kuhlenbäumer G, Bevan S, Jonsdottir T, Bjarnason H, Saemundsdottir J, Palsson S, Arnar DO, Holm H, Thorgeirsson G, Valdimarsson EM, Sveinbjörnsdottir S, Gieger C, Berger K, Wichmann HE, Hillert J, Markus H, Gulcher JR, Ringelstein EB, Kong A, Dichgans M, Gudbjartsson DF, Thorsteinsdottir U, Stefansson K.

Risk variants for atrial fibrillation on chromosome 4q25 associate with ischemic stroke. *Ann Neurol.* 2008;64(4): 402-9

Rösser N, Heuschmann P, Wersching H, Breitenstein C, Knecht S, Flöel A.

Levodopa improves procedural motor learning in chronic stroke patients. *Arch Phys Med Rehabil.* 2008;89(9): 1633-41

Weikert C, Stefan N, Schulze MB, Pischon T, Berger K, Joost HG, Häring HU, Boeing H, Fritsche A.

Plasma fetuin-a levels and the risk of myocardial infarction and ischemic stroke. *Circulation.* 2008;118(24): 2555-62

Frey UH, Lieb W, Erdmann J, Savidou D, Heusch G, Leineweber K, Jakob H, Hense HW, Löwel H, Brockmeyer NH, Schunkert H, Siffert W.

Characterization of the GNAQ promoter and association of increased Gq expression with cardiac hypertrophy in humans. *Eur Heart J.* 2008;29(7): 888-97

Knecht S, Oelschlager C, Duning T, Lohmann H, Albers J, Stehling C, Heindel W, Breithardt G, Berger K, Ringelstein EB, Kirchhof P, Wersching H.

Atrial fibrillation in stroke-free patients is associated with memory impairment and hippocampal atrophy. *Eur Heart J.* 2008;29(17): 2125-2132

Hense HW, Koesters E, Wellmann J, Meisinger C, Völzke H, Keil U.

Evaluation of a recalibrated Systematic Coronary Risk Evaluation cardiovascular risk chart: results from Systematic Coronary Risk Evaluation Germany. *Eur J Cardiovasc Prev Rehabil.* 2008;15(4): 409-15

Prugger C, Wellmann J, Heidrich J, Brand-Herrmann SM, Keil U.

Cardiovascular risk factors and mortality in patients with coronary heart disease. *Eur J Epidemiol.* 2008;23(11): 731-7

Störk S, Hense HW, Zentgraf C, Uebelacker I, Jahns R, Ertl G, Angermann CE.

Pharmacotherapy according to treatment guidelines is associated with lower mortality in a community-based sample of patients with chronic heart failure: a prospective cohort study. *Eur J Heart Fail.* 2008;10(12): 1236-45

Weinmayr G, Forastiere F, Weiland SK, Rzehak P, Abramidze T, Annesi-Maesano I, Björkstén B, Brunekreef B, Büchele G, Cookson WO, von Mutius E, Pistelli R, Strachan DP, ISAAC Phase Two Study Group.

International variation in prevalence of rhinitis and its relationship with sensitisation to perennial and seasonal allergens. *Eur Respir J.* 2008;32(5): 1250-61

Knecht S, Wersching H, Lohmann H, Bruchmann M, Duning T, Dziewas R, Berger K, Ringelstein EB.

High-normal blood pressure is associated with poor cognitive performance. *Hypertension.* 2008;51(3): 663-8

Azevedo A, Bettencourt P, Alvelos M, Martins E, Abreu-Lima C, Hense HW, Barros H.

Health-related quality of life and stages of heart failure. *Int J Cardiol.* 2008;129(2): 238-44

Anderson HR, Gupta R, Kapetanakis V, Asher MI, Clayton T, Robertson CF, Strachan DP, ISAAC Steering Committee.

International correlations between indicators of prevalence, hospital admissions and mortality for asthma in children. *Int J Epidemiol.* 2008;37(3): 573-82

Flohr C, Weiland SK, Weinmayr G, Björkstén B, Bråbäck L, Brunekreef B, Büchele G, Clausen M, Cookson WO, von Mutius E, Strachan DP, Williams HC, ISAAC Phase Two Study Group.

The role of atopic sensitization in flexural eczema: findings from the International Study of Asthma and Allergies in Childhood Phase Two. *J Allergy Clin Immunol.* 2008;121(1): 141-147.e4

Williams H, Stewart A, von Mutius E, Cookson W, Anderson HR, International Study of Asthma and Allergies in Childhood (ISAAC) Phase One and Three Study Groups.

Is eczema really on the increase worldwide? *J Allergy Clin Immunol.* 2008;121(4): 947-54.e15

Buchner S, Muscholl M, Debl K, Hense HW, Döring A, Stritzke J, Schunkert H, Jougasaki M, Burnett JC, Riegger GA, Luchner A.

Left atrial size by planimetry is superior to M-mode diameter: biochemical calibration by atrial and brain natriuretic peptide. *J Am Soc Echocardiogr.* 2008;21(4): 380-5

Weikert C, Westphal S, Berger K, Dierkes J, Möhlig M, Spranger J, Rimm EB, Willich SN, Boeing H, Pischon T.

Plasma resistin levels and risk of myocardial infarction and ischemic stroke. *J Clin Endocrinol Metab.* 2008;93(7): 2647-53

Markus MR, Stritzke J, Lieb W, Mayer B, Luchner A, Döring A, Keil U, Hense HW, Schunkert H.

Implications of persistent prehypertension for ageing-related changes in left ventricular geometry and function: the MONICA/KORA Augsburg study. *J Hypertens.* 2008;26(10): 2040-9

Happe S, Vennemann M, Evers S, Berger K.

Treatment wish of individuals with known and unknown restless legs syndrome in the community. *J Neurol.* 2008;255(9): 1365-71

Vennemann MM, Hummel T, Berger K.

The association between smoking and smell and taste impairment in the general population. *J Neurol.* 2008;255(8): 1121-6

Beasley R, Clayton T, Crane J, von Mutius E, Lai CK, Montefort S, Stewart A, ISAAC Phase Three Study Group.

Association between paracetamol use in infancy and childhood, and risk of asthma, rhinoconjunctivitis, and eczema in children aged 6-7 years: analysis from Phase Three of the ISAAC programme. *Lancet.* 2008;372(9643): 1039-48

Greve B, Hoffmann P, Vonthein R, Kun J, Lell B, Mycko MP, Selmaj KW, Berger K, Weissert R, Kremsner PG.

NCF1 gene and pseudogene pattern: association with parasitic infection and autoimmunity. *Malar J.* 2008;7: 251

Greve B, Simonenko R, Illes Z, Peterfalvi A, Hamdi N, Mycko MP, Selmaj KW, Rozsa C, Rajczy K, Bauer P, Berger K, Weissert R.

Multiple sclerosis and the CTLA4 autoimmunity polymorphism CT60: no association in patients from Germany, Hungary and Poland. *Mult Scler.* 2008;14(2): 153-8

Rettig K, Trenkwalder C, Berger K.

Prevalence of restless legs syndrome in primary care - a study in 311 German medical practices. *NERVENHEILKUNDE.* 2008;27(4): 334-338

Baune BT, Ponath G, Golledge J, Varga G, Arolt V, Rothermundt M, Berger K.

Association between IL-8 cytokine and cognitive performance in an elderly general population--the MEMO-Study. *Neurobiol Aging.* 2008;29(6): 937-44

Flöel A, Witte AV, Lohmann H, Wersching H, Ringelstein EB, Berger K, Knecht S.

Lifestyle and memory in the elderly. *Neuroepidemiology.* 2008;31(1): 39-47

Baune BT, Hohoff C, Berger K, Neumann A, Mortensen S, Roehrs T, Deckert J, Arolt V, Domschke K.

Association of the COMT val158met variant with antidepressant treatment response in major depression. *Neuropsychopharmacology.* 2008;33(4): 924-32

de Vocht F, Vermeulen R, Burstyn I, Sobala W, Dost A, Taege D, Bergendorf U, Straif K, Swuste P, Kromhout H, EU-EXASRUB consortium.

Exposure to inhalable dust and its cyclohexane soluble fraction since the 1970s in the rubber manufacturing industry in the European Union. *Occup Environ Med.* 2008;65(6): 384-91

Neuner B, Wellmann J, Dasch B, Dietzel M, Farwick A, Stoll M, Pauleikhoff D, Hense HW.

LOC387715, smoking and their prognostic impact on visual functional status in age-related macular degeneration-The Muenster Aging and Retina Study (MARS) cohort. *Ophthalmic Epidemiol.* 2008;15(3): 148-54

Baune BT, Caniato RN, Garcia-Alcaraz MA, Berger K.

Combined effects of major depression, pain and somatic disorders on general functioning in the general adult population. *Pain.* 2008;138(2): 310-7

Björkstén B, Clayton T, Ellwood P, Stewart A, Strachan D, ISAAC Phase III Study Group.

Worldwide time trends for symptoms of rhinitis and conjunctivitis: Phase III of the International Study of Asthma and Allergies in Childhood. *Pediatr Allergy Immunol.* 2008;19(2): 110-24

Rösser N, Berger K, Vomhof P, Knecht S, Breitenstein C, Flöel A.

Lack of improvement in odor identification by levodopa in humans. *Physiol Behav.* 2008;93(4-5): 1024-9

Richter D, Berger K, Reker T.

[Are mental disorders on the increase? A systematic review] *Psychiatr Prax.* 2008;35(7): 321-30

Baune BT, Ponath G, Rothermundt M, Riess O, Funke H, Berger K.

Association between genetic variants of IL-1beta, IL-6 and TNF-alpha cytokines and cognitive performance in the elderly general population of the MEMO-study. *Psychoneuroendocrinology.* 2008;33(1): 68-76

Azevedo A, Bettencourt P, Friões F, Alvelos M, Abreu-Lima C, Hense HW, Barros H.

Depressive symptoms and heart failure stages. Psychosomatics. 2008;49(1): 42-8

Minnerup J, Heidrich J, Wellmann J, Rogalewski A, Schneider A, Schäbitz WR.  
Meta-analysis of the efficacy of granulocyte-colony stimulating factor in animal models of focal cerebral ischemia. Stroke. 2008;39(6): 1855-61

Kraywinkel K, Lehnert M, Semjonow A, Hense HW.  
[Epidemiology of prostate cancer: recent results from the Epidemiological Cancer Register of the District of Münster (Germany)] Urologe. 2008;47(7): 853-9