

[2010]

Heuschmann PU, Busse O, Wagner M, Endres M, Villringer A, Rother J, Kolominsky-Rabas PL, Berger K, Deut Schlaganfall Gesell; Stiftung Deutsch Schlaganfall. Frequency and Care of Stroke in Germany. AKTUEL NEUROL. 2010;37(7): 333-340

van den Bussche H, Berger K, Kemper C, Barzel A, Glaeske G, Koller D. Incidence, Relapse, Nursing Care Dependency and Mortality of Stroke in Germany A Secondary Analysis of Statutory Insurance Claims Data. AKTUEL NEUROL. 2010;37(3): 131-135

Baune BT, Neuhauser H, Ellert U, Berger K. The role of the inflammatory markers ferritin, transferrin and fibrinogen in the relationship between major depression and cardiovascular disorders - The German Health Interview and Examination Survey. Acta Psychiatr Scand. 2010;121(2): 135-42

Weinmayr G, Genuneit J, Nagel G, Björkstén B, van Hage M, Priftanji A, Cooper P, Rijkjäv MA, von Mutius E, Tsanakas J, Forastiere F, Doekes G, Garrido JB, Suarez-Varela MM, Bråbäck L, Strachan DP, ISAAC Phase Two Study Group. International variations in associations of allergic markers and diseases in children: ISAAC Phase Two. Allergy. 2010;65(6): 766-75

Berger K, Evers S. Migraine with aura and the risk of increased mortality. BMJ. 2010;341: c4410

Diederichs C, Pohlmann R, Berger K. [The frequency of older people living in nursing homes: results from a population-based study in Dortmund]. Bundesgesundheitsblatt Gesundheitsforschung, Gesundheitsschutz. 2010;53(5): 504-9

Straube A, Pfaffenrath V, Ladwig KH, Meisinger C, Hoffmann W, Fendrich K, Vennemann M, Berger K. Prevalence of chronic migraine and medication overuse headache in Germany--the German DMKG headache study. Cephalalgia. 2010;30(2): 207-13

Kuhlenbaeumer G, Hüge A, Berger K, Kessler C, Voelzke H, Funke H, Stoegbauer F, Stoll M, Ringelstein EB. Genetic variants in the C-reactive protein gene are associated with microangiopathic ischemic stroke. Cerebrovasc Dis. 2010;30(5): 476-82

Meves SH, Diehm C, Berger K, Pittrow D, Trampisch HJ, Burghaus I, Tepohl G, Allenberg JR, Endres HG, Schwertfeger M, Darius H, Haberl RL, getABI Study Group. Peripheral arterial disease as an independent predictor for excess stroke morbidity and mortality in primary-care patients: 5-year results of the getABI study. Cerebrovasc Dis. 2010;29(6): 546-54

De Bacquer D, Dallongeville J, Heidrich J, Kotseva K, Reiner Z, Gaita D, Prugger C, Wellmann J, Wood D, De Backer G, Keil U, EUROASPIRE III Study Group. Management of overweight and obese patients with coronary heart disease across Europe. Eur J Cardiovasc Prev Rehabil. 2010;17(4): 447-54

Kotseva K, Wood D, De Backer G, De Bacquer D, Pyörälä K, Reiner Z, Keil U, EUROASPIRE Study Group.

EUROASPIRE III. Management of cardiovascular risk factors in asymptomatic high-risk patients in general practice: cross-sectional survey in 12 European countries. Eur J Cardiovasc Prev Rehabil. 2010;17(5): 530-40

Minnerup J, Wersching H, Schäbitz WR.

EPO for stroke therapy - Is there a future for further clinical development? Exp Transl Stroke Med. 2010;2(1): 10

Neuner B, Berger K.

[Individual social factors and their association with environmental socioeconomic factors--a descriptive small-area analysis in the city of Dortmund]. Gesundheitswesen. 2010;72(11): 797-803

Wieser T, Dresler K, Evers S, Gaul C, König D, Hölzl D, Berger K, Nyholt D, Deufel T. No influence of 5-HTTLPR gene polymorphism on migraine symptomatology, comorbid depression, and chronification. Headache. 2010;50(3): 420-30

Mehrhof F, Löffler M, Gelbrich G, Ozcelik C, Posch M, Hense HW, Keil U, Scheffold T, Schunkert H, Angermann C, Ertl G, Jahns R, Pieske B, Wachter R, Edelmann F, Wollert KC, Maisch B, Pankuweit S, Erbel R, Neumann T, Herzog W, Katus H, Müller-Tasch T, Zugck C, Düngen HD, Regitz-Zagrosek V, Lehmkühl E, Störk S, Siebert U, Wasem J, Neumann A, Göhler A, Anker SD, Köhler F, Möckel M, Osterziel KJ, Dietz R, Rauchhaus M, Competence Network Heart Failure.

A network against failing hearts--introducing the German "Competence Network Heart Failure". Int J Cardiol. 2010;145(1): 135-8

Domschke K, Lawford B, Laje G, Berger K, Young R, Morris P, Deckert J, Arolt V, McMahon FJ, Baune BT.

Brain-derived neurotrophic factor (BDNF) gene: no major impact on antidepressant treatment response. Int J Neuropsychopharmacol. 2010;13(1): 93-101

Ellwood P, Asher MI, Stewart AW, ISAAC Phase III Study Group.

The impact of the method of consent on response rates in the ISAAC time trends study. Int J Tuberc Lung Dis. 2010;14(8): 1059-65

Farwick A, Wellmann J, Stoll M, Pauleikhoff D, Hense HW.

Susceptibility genes and progression in age-related maculopathy: a study of single eyes. Invest Ophthalmol Vis Sci. 2010;51(2): 731-6

Assimes TL, Hólm H, Kathiresan S, Reilly MP, Thorleifsson G, Voight BF, Erdmann J, Willenborg C, Vaidya D, Xie C, Patterson CC, Morgan TM, Burnett MS, Li M, Hlatky MA, Knowles JW, Thompson JR, Absher D, Iribarren C, Go A, Fortmann SP, Sidney S, Risch N, Tang H, Myers RM, Berger K, Stoll M, Shah SH, Thorleifsson G, Andersen K, Havulinna AS, Herrera JE, Faraday N, Kim Y, Kral BG, Mathias RA, Ruczinski I, Sukitipat B, Wilson AF, Yanek LR, Becker LC, Linsel-Nitschke P, Lieb W, König IR, Hengstenberg C, Fischer M, Stark K, Reinhard W, Winogradow J, Grassl M, Grosshennig A, Preuss M, Schreiber S, Wichmann HE, Meisinger C, Yee J, Friedlander Y, Do R, Meigs JB, Williams G, Nathan DM, MacRae CA, Qu L, Wilensky RL, Matthai WH, Qasim AN, Hakonarson H, Pichard AD, Kent KM, Satler L, Lindsay JM, Waksman R, Knouff CW, Waterworth DM, Walker MC, Mooser VE, Marrugat J, Lucas G, Subirana I, Sala J, Ramos R, Martinelli N, Olivieri O, Trabetti E, Malerba G, Pignatti PF, Guiducci C, Mirel D, Parkin M, Hirschhorn JN, Asselta R, Duga S, Musunuru K, Daly MJ, Purcell S, Eifert S, Braund PS, Wright BJ, Balmforth AJ, Ball SG, Myocardial Infarction Genetics Consortium, Wellcome Trust Case Control Consortium, Cardiogenics, Ouwehand WH, Deloukas P, Scholz M, Cambien F, Hogue A, Scheffold T, Salomaa V, Girelli D, Granger CB, Peltonen L, McKeown PP, Altshuler D, Melander O,

Devaney JM, Epstein SE, Rader DJ, Elosua R, Engert JC, Anand SS, Hall AS, Ziegler A, O'Donnell CJ, Spertus JA, Siscovick D, Schwartz SM, Becker D, Thorsteinsdottir U, Stefansson K, Schunkert H, Samani NJ, Quertermous T.
Lack of association between the Trp719Arg polymorphism in kinesin-like protein-6 and coronary artery disease in 19 case-control studies. *J Am Coll Cardiol*. 2010;56(19): 1552-63

Markus MR, Stritzke J, Siewert U, Lieb W, Luchner A, Döring A, Keil U, Hense HW, Schunkert H, MONICA/KORA Investigators.
Variation in body composition determines long-term blood pressure changes in pre-hypertension: the MONICA/KORA (Monitoring Trends and Determinants on Cardiovascular Diseases/Cooperative Research in the Region of Augsburg) cohort study. *J Am Coll Cardiol*. 2010;56(1): 65-76

Minnerup J, Wersching H, Diederich K, Schilling M, Ringelstein EB, Wellmann J, Schäbitz WR.
Methodological quality of preclinical stroke studies is not required for publication in high-impact journals. *J Cereb Blood Flow Metab*. 2010;30(9): 1619-24

Gehrmann U, Kuss O, Wellmann J, Bender R.
Logistic regression was preferred to estimate risk differences and numbers needed to be exposed adjusted for covariates. *J Clin Epidemiol*. 2010;63(11): 1223-31

Schoenfeld H, Pruss A, Keller M, Schuster M, Meinck K, Neuner B, von Heymann C.
Lyophilised plasma: evaluation of clotting factor activity over 6 days after reconstitution for transfusion. *J Clin Pathol*. 2010;63(8): 726-30

Jung A, Hüge A, Kühlenbäumer G, Kempt S, Seehafer T, Evers S, Berger K, Marziniak M.
Genetic TPH2 variants and the susceptibility for migraine: association of a TPH2 haplotype with migraine without aura. *J Neural Transm*. 2010;117(11): 1253-60

Pham TD, Elfiqi HZ, Knecht S, Wersching H, Baune BT, Berger K.
Structural simplicity of the brain. *J Neurosci Methods*. 2010;188(1): 113-26

Lindinger A, Schwedler G, Hense HW.
Prevalence of congenital heart defects in newborns in Germany: Results of the first registration year of the PAN Study (July 2006 to June 2007). *Klin Padiatr*. 2010;222(5): 321-6

Cheema B, Abas H, Smith B, O'Sullivan AJ, Chan M, Patwardhan A, Kelly J, Gillin A, Pang G, Lloyd B, Berger K, Baune BT, Singh MF.
Investigation of skeletal muscle quantity and quality in end-stage renal disease *Nephrology (Carlton)*. 2010;15(4): 454-463

Pohl M, Berger K, Ketter G, Krusch C, Pause M, Puschendorf W, Schaupp M, Schleep J, Spranger M, Steube D, Scheidtmann K, Mehrholz J.
[Long-term course of patients in neurological rehabilitation Phase B : Results of the 6-year follow-up in a multicenter study.] *Nervenarzt*. 2010;82(6): 753-763

Wersching H, Dünning T, Lohmann H, Mohammadi S, Stehling C, Fobker M, Conty M, Minnerup J, Ringelstein EB, Berger K, Deppe M, Knecht S.
Serum C-reactive protein is linked to cerebral microstructural integrity and cognitive function. *Neurology*. 2010;74(13): 1022-9

Anderson HR, Ruggles R, Pandey KD, Kapetanakis V, Brunekreef B, Lai CK, Strachan DP, Weiland SK, ISAAC Phase One Study Group.

Ambient particulate pollution and the world-wide prevalence of asthma, rhinoconjunctivitis and eczema in children: Phase One of the International Study of Asthma and Allergies in Childhood (ISAAC). *Occup Environ Med.* 2010;67(5): 293-300

Walters AS, Moussouttas M, Siddiqui F, Silveira DC, Fuentes K, Wang L, Berger K. Prevalence of stroke in Restless Legs Syndrome: Initial Results Point to the Need for More Sophisticated Studies. *Open Neurol J.* 2010;4: 73-7

Dietzel M, Farwick A, Hense HW. [Genetic and risk factors for exudative AMD]. *Ophthalmologe.* 2010;107(12): 1103-8

Bücheler G, Genuneit J, Weinmayr G, Björkstén B, Gehring U, von Mutius E, Priftanji A, Stein RT, Addo-Yobo EO, Priftis KN, Shah JR, Forastiere F, Svabe V, Crane J, Nystad W, García-Marcos L, Saraçlar Y, El-Sharif N, Strachan DP, ISAAC Phase Two Study Group. International variations in bronchial responsiveness in children: findings from ISAAC phase two. *Pediatr Pulmonol.* 2010;45(8): 796-806

Asher MI, Stewart AW, Mallol J, Montefort S, Lai CK, Ait-Khaled N, Odhiambo J, ISAAC Phase One Study Group. Which population level environmental factors are associated with asthma, rhinoconjunctivitis and eczema? Review of the ecological analyses of ISAAC Phase One. *Respir Res.* 2010;11: 8

Berger K. What is clinically significant RLS and who decides about its treatment? *Sleep Med.* 2010;11(1): 9-10

Rist PM, Berger K, Buring JE, Kase CS, Gaziano JM, Kurth T. Alcohol consumption and functional outcome after stroke in men. *Stroke.* 2010;41(1): 141-6

Schoenfeld H, Perka C, Ziemer S, Huebner R, Schink T, Neuner B, Spies C. The perioperative von Willebrand factor activity and factor VIII levels among alcohol use disorder patients undergoing total knee or hip replacement. *Subst Use Misuse.* 2010;45(7-8): 1216-29

Nagel G, Weinmayr G, Kleiner A, Garcia-Marcos L, Strachan DP, ISAAC Phase Two Study Group. Effect of diet on asthma and allergic sensitisation in the International Study on Allergies and Asthma in Childhood (ISAAC) Phase Two. *Thorax.* 2010;65(6): 516-22

Schoenfeld H, Pretzel KJ, von Heymann C, Neuner B, Kalus U, Kiesewetter H, Pruss A. Validation of a hospital-laboratory workstation for immunohematologic methods. *Transfusion.* 2010;50(1): 26-31