

1997

1. Hirsch JR, Cermak R, Forssmann W-G, Kleta R, Kruhøffer M, Kuhn M, Schafer JA, Sun D, Schlatter E. Effects of sodium nitroprusside in the rat cortical collecting duct (CCD) are independent of the NO-pathway. *Kidney Int*, 1997;51:473-476
2. Schlatter E, Cermak R, Hirsch J, Kuhn M, Kleta R, Forssmann W-G, Sun D, Schafer JA. cGMP activating peptides do not regulate electrogenic electrolyte transport in principal cells of rat CCD. *Am J Physiol Renal Physiol*, 1997;271:F1158-1165
3. Ankorina I, Haxelmans S, Schlatter E. Functional evidence for the regulation of cytosolic Ca^{2+} activity via V1-receptors and β -adrenoceptors in rat CCD. *Cell Calcium*, 1997;21:163-171
4. Hirsch JR, Schlatter E. Ca^{2+} -dependent K^{+} channels in the cortical collecting duct of rat. *Wien Klin Wochenschr*, 1997;109:485-488 [IF 1997: 0,411]
5. Schlatter E, Ankorina I, Haxelmans S, Hohage H. Moxonidine inhibits $\text{Na}^{+}/\text{H}^{+}$ exchange in proximal tubule cells and cortical collecting duct. *Kidney Int*, 1997;52:454-459
6. Schafer JA, Watkins ML, Li L, Herter P, Haxelmans S, Schlatter E. A simplified method for isolation of large numbers of defined nephron segments. *Am J Physiol Renal Physiol*; 1997;273:F650-F657
7. Hohage H, Schlatter E, Greven J. Effects of moxonidine and clonidine on renal function and blood pressure in anesthetized rats. *Clin Nephrol*, 1997;47:316-324
8. Kleta R, Lennert T, Versmold H, Schlatter E. Diadenosin-Polyphosphate und renale Mesangialzellen. *Anonymous Jahrbuch UKBF der Freien Universität Berlin*. 1997;2:224-225
9. Stachon A, Hohage H, Feidt C, Kleta R, Schlatter E. Characterization of organic cation transport across the apical membrane of proximal tubular cells with the fluorescent dye 4-Di-1-ASP. *Cell Physiol Biochem*, 1997;7:264-274
10. Schlatter E, Ankorina-Stark I, Cermak R, Haxelmans S, Kleta R, Hirsch JR. Cell shrinkage activates a cation conductance in principal cells of rat cortical collecting duct. *Cell Physiol Biochem*, 1997;7:321-332
11. Ankorina-Stark I, Haxelmans S, Hirsch JR, Lohrmann E, Schlatter E. Ca^{2+} entry in isolated rat cortical collecting duct is pH- and voltage-sensitive. Different pathways for basal and swelling-induced Ca^{2+} entry. *Cell Physiol Biochem*, 1997;7:333-344
12. Ankorina-Stark I, Haxelmans S, Schlatter E. Receptors for bradykinin and prostaglandin E2 coupled to Ca^{2+} signaling in rat cortical collecting duct. *Cell Calcium*, 1997;22:269-275
13. Hohage H, Hess K, Jahl C, Greven J, Schlatter E. Renal and blood pressure effects of moxonidine and clonidine in spontaneously hypertensive rats. *Clin Nephrol*, 1997;48:346-352